

Scattering induced current in a tight-binding band

L. Bruneau*

Dépt. de Mathématiques and UMR 8088, CNRS

Université de Cergy-Pontoise[†]

and

S. De Bièvre[‡]

Lab. P. Painlevé, CNRS, UMR 8524 and UFR de Mathématiques

Université Lille 1, Sciences et Technologies[§]

Équipe-Projet SIMPAF, Centre de Recherche INRIA Futurs[¶]

and

C.-A. Pillet^{||}

Centre de Physique Théorique**

Université du Sud Toulon-Var^{††}

January 18, 2011

Abstract. In the single band tight-binding approximation, we consider the transport properties of an electron subject to a homogeneous static electric field. We show that repeated interactions of the electron with two-level systems in thermal equilibrium suppress the Bloch oscillations and induce a steady current, the statistical properties of which we study.

*laurent.bruneau@u-cergy.fr

†95000 Cergy-Pontoise, France

‡Stephan.De-Bievre@math.univ-lille1.fr

§F-59655 V'd'Ascq Cedex, France

¶Parc Scient. de la Haute Borne, 40, avenue Halley B.P. 70478, F-59658 V'd'Ascq cedex, France

||pillet@univ-tln.fr

**UMR 6207: Univ. de Provence, Univ. de la Méditerranée, Univ. de Toulon et CNRS, FRUMAM

††B.P. 20132, 83957 La Garde Cedex, France

1 Introduction

Let us consider the motion of an electron in the single band tight-binding approximation. It is well known that in the absence of a dc electric field, the electron moves ballistically whereas, when a dc electric field is present, Bloch oscillations prevent a current from being set up in the system. It is furthermore expected that if the electron is in contact with a thermal environment, the resulting scattering mechanisms will suppress the Bloch oscillations and lead to a steady current. This phenomenon can for example be described within a semi-classical picture of the motion combined with the relaxation-time approximation (see [AM] for example). Alternatively, in open systems theory, one describes the thermal environment and its coupling to the electron with a global Hamiltonian, and traces out the environment variables to obtain, under suitable additional conditions such as weak coupling and appropriate scalings, effective dynamical equations for the electron alone. In treatments of this type, the environment is often described by a set of oscillators (see for example [FZ]) or more generally by a free Bose field. For a review of various approaches to transport theory we refer to [B]. There has recently been intensified interest in obtaining rigorous results along those lines [DFP, DF, CDM].

We present here a simple, explicitly solvable, fully quantum mechanical and fully Hamiltonian model of such a particle-environment system within the repeated interaction scheme, in which the environment is described by a chain of two-level atoms. We show that a dc current is created due to the interaction of the particle with its environment. In addition to drifting in the direction of the applied field, the electron diffuses around its mean position. We give a full analysis of the probability distribution for the position of the particle in the large time regime (Theorem 3.1). We then use a repeated measurement scheme to describe the increments of the position and energy observables between time 0 and time t (Theorems 3.3 and 3.2).

The rest of the paper is organized as follows. In Section 2 we give a detailed description of the model under consideration. Our main results are stated precisely in Section 3. We describe the effect on the particle of the interaction with a single atom in Section 4, and the main properties of the repeated interaction dynamics in Section 5. The proofs of the main theorems are provided in Sections 6 and 7.

2 Description of the model

We consider a spinless particle on the one-dimensional lattice \mathbb{Z} and submitted to a constant external force $F \geq 0$. The quantum Hilbert space and Hamiltonian of the particle are

$$\mathcal{H}_p = \ell^2(\mathbb{Z}), \quad H_p = -\Delta - FX, \quad (2.1)$$

where Δ is the usual discrete nearest neighbor Laplacian and X the lattice position operator

$$-\Delta = \sum_{x \in \mathbb{Z}} (2|x\rangle\langle x| - |x+1\rangle\langle x| - |x\rangle\langle x+1|), \quad X = \sum_{x \in \mathbb{Z}} x|x\rangle\langle x|.$$

Identifying \mathcal{H}_p with $L^2(\mathbb{T}^1, d\xi)$ via the discrete Fourier transform, we have

$$-\Delta = 2(1 - \cos \xi), \quad X = i\partial_\xi.$$

Here $\mathbb{T}^1 \simeq [0, 2\pi[$ is the first Brillouin zone and ξ the crystal momentum. Defining the translation operator

$$T = \sum_{x \in \mathbb{Z}} |x+1\rangle\langle x| = e^{-i\xi},$$

we can write $H_p = 2 - T - T^* - FX$. We note for later reference that

$$[X, T] = T, \quad [X, T^*] = -T^*, \quad [H_p, T] = -FT, \quad [H_p, T^*] = FT^*. \quad (2.2)$$

When $F = 0$, H_p has a single band of absolutely continuous spectrum, $\text{sp}(H_p) = [0, 4]$, and the motion of the particle is described by

$$\begin{aligned} T(t) &= e^{itH_p} T e^{-itH_p} = T, \\ X(t) &= e^{itH_p} X e^{-itH_p} = X + i(T - T^*)t, \end{aligned}$$

showing its ballistic nature.

When $F \neq 0$, we set $G = 2F^{-1} \sin \xi = iF^{-1}(T - T^*)$. The commutation relations (2.2) yield

$$H_p = e^{iG}(2 - FX)e^{-iG}, \quad (2.3)$$

from which it follows that H_p has discrete spectrum, $\text{sp}(H_p) = 2 - F\mathbb{Z}$. This is the well-known Wannier-Stark ladder. In the momentum representation, the normalized eigenvector ψ_k to the eigenvalue $E_k = 2 - Fk$ is given by

$$\widehat{\psi}_k(\xi) = \frac{1}{\sqrt{2\pi}} e^{i(2F^{-1} \sin(\xi) - k\xi)}. \quad (2.4)$$

In the position representation, we therefore have

$$\psi_k(x) = \int_0^{2\pi} e^{i(2F^{-1} \sin(\xi) - (k-x)\xi)} \frac{d\xi}{2\pi} = J_{k-x} \left(\frac{2}{F} \right),$$

where the J_ν are Bessel functions. From their asymptotic behavior for large ν (see e.g. Formula (10.19.1) in [OLBC]) we infer that

$$\psi_k(x) \sim \frac{1}{\sqrt{2\pi|k-x|}} \left(\frac{e}{F|k-x|} \right)^{|k-x|} \quad \text{for } |k-x| \rightarrow \infty,$$

which shows that $\psi_k(x)$ is sharply localized around $x = k$. The motion of the particle, described by

$$\left. \begin{aligned} T(t) &= e^{itH_p} T e^{-itH_p} = e^{-itF} T, \\ X(t) &= e^{itH_p} X e^{-itH_p} = X + \frac{4}{F} \sin\left(\frac{Ft}{2}\right) \sin\left(\xi + \frac{Ft}{2}\right), \end{aligned} \right\} \quad (2.5)$$

is now confined by Bloch oscillations.

In what follows, we let the particle interact with one or many 2-level atoms, each of which has a quantum Hilbert space $\mathcal{H}_a = \mathbb{C}^2$ which we identify with $\Gamma_-(\mathbb{C})$, the fermionic Fock space over \mathbb{C} . The atomic Hamiltonian is given by

$$H_a = d\Gamma(E) = Eb^*b,$$

where $E \geq 0$ is the Bohr frequency of the atom and b^* , b are the usual Fermi creation and annihilation operators. The later satisfy the canonical anti-commutation relations

$$b^*b + bb^* = I, \quad b^2 = b^{*2} = 0. \quad (2.6)$$

The initial state of the two-level atoms will be their equilibrium state at inverse temperature β described by the density matrix

$$\rho_\beta = Z_\beta^{-1} e^{-\beta H_a}, \quad Z_\beta = \text{Tr}(e^{-\beta H_a}) = 1 + e^{-\beta E}. \quad (2.7)$$

The interaction between the particle and the two-level atom is chosen so that its effect is to give a right or left kick to the particle, depending on whether the atom is in its ground state or in its excited state. More precisely, we set

$$V = \sum_{x \in \mathbb{Z}} (|x+1\rangle\langle x| \otimes b^* + |x\rangle\langle x+1| \otimes b) = Tb^* + T^*b. \quad (2.8)$$

To understand this interaction, note that when $F > 0$, the translation operator T can be thought of as a lowering operator for the particle. Indeed, from (2.4) one finds

$$T\psi_k = \psi_{k+1}. \quad (2.9)$$

Similarly, T^* acts as a raising operator. As a result, V describes an exchange of energy between the two-level system and the particle. We point out that this interaction is very similar to the one which appears in the Jaynes-Cummings Hamiltonian where a two level atom interacts with one mode of the electromagnetic field of a cavity (through its electric dipole moment and in the rotating wave approximation), see e.g. [CDG, Du]. Thermalization of the field through repeated interaction with two-level

atoms was proven for the Jaynes-Cummings Hamiltonian in [BP]. The model treated here is very similar to the one studied in [BP], except that the spectrum of H_p , contrary to the spectrum of the mode of the electromagnetic field, is not bounded from below. As a result, the system we treat here has no invariant state, as we shall see below.

The full Hamiltonian of the particle interacting with a single two-level system acts on the Hilbert space $\mathcal{H}_p \otimes \mathcal{H}_a$ and is given by

$$H = H_p + H_a + \lambda V, \quad (2.10)$$

where $\lambda \in \mathbb{R}$ is a coupling constant. As the more explicit formula

$$H = 2 - T(1 - \lambda b^*) - T^*(1 - \lambda b) - FX + Eb^*b,$$

shows, one can also interpret the coupling to the two-level system as altering the hopping matrix elements of the original Hamiltonian. The operator H is easily diagonalized by noticing that it commutes with the “number operator”

$$N = -e^{iG} X e^{-iG} + b^*b = \frac{H_p - 2}{F} + \frac{H_a}{E}, \quad (2.11)$$

which has a two-dimensional eigenspace to each of its eigenvalue $n \in \mathbb{Z}$. In particular, if $E = F$ then the energy $H_p + H_a$ is preserved by the full dynamics (which will be computed in Section 4).

We now turn to the description of the repeated interaction dynamics (see e.g. [AP, BJM, BP]). We let the particle interact successively, each time during a fixed period $\tau > 0$, with the elements of a sequence of atoms, *i.e.*, during the time interval $[(n-1)\tau, n\tau[$, the particle interacts with the n -th atom and with none of the others. The Hilbert space of the atomic reservoir is

$$\mathcal{H}_{\text{env}} = \bigotimes_{n=1}^M \mathcal{H}_{a,n},$$

where M is the number of atoms and each $\mathcal{H}_{a,n}$ is a copy of \mathcal{H}_a . The Hilbert space of the joint particle+reservoir system is $\mathcal{H} = \mathcal{H}_p \otimes \mathcal{H}_{\text{env}}$. The full unitary evolution $U(t, s)$ of the system is thus described by the Schrödinger equation

$$i\partial_t U(t, s) = H(t)U(t, s), \quad U(s, s) = I, \quad t, s \in [0, M\tau], \quad (2.12)$$

with time-dependent Hamiltonian

$$H(t) = H_p + \sum_{n=1}^M H_{a,n} + \lambda \sum_{n=1}^M \chi_n(t) (Tb_n^* + T^*b_n), \quad (2.13)$$

where χ_n is the characteristic function of the interval $[(n-1)\tau, n\tau[$ and $H_{a,n}$, b_n , b_n^* are the Hamiltonian, annihilation and creation operators of the n -th atom. We will use the following notation

$$H_n = H_p + H_{a,n} + \lambda(Tb_n^* + T^*b_n), \quad \tilde{H}_n = H_n + \sum_{\substack{1 \leq k \leq M \\ k \neq n}} H_{a,k}. \quad (2.14)$$

Note that $H(t) = \tilde{H}_n$ when $t \in [(n-1)\tau, n\tau[$.

We denote by $\mathcal{B}^1(\mathcal{H}_p)$ the Banach space of trace class operators on the Hilbert space \mathcal{H}_p . Given any density matrix for the particle $\rho_p \in \mathcal{B}^1(\mathcal{H}_p)$, $\rho_p \geq 0$, $\text{Tr}\rho_p = 1$, we set the initial state of the joint system to

$$\rho_0 = \rho_p \otimes \rho_\beta^{\otimes M}, \quad \rho_\beta^{\otimes M} = \bigotimes_{n=1}^M \rho_\beta.$$

After $n \leq M$ interactions, this state evolves into $\rho_{n\tau} = U(n\tau, 0)\rho_0U(n\tau, 0)^*$. To obtain the density matrix $\rho_{p,n\tau}$ of the particle after these n interactions we take the partial trace over the environment,

$$\rho_{p,n\tau} = \text{Tr}_{\mathcal{H}_{\text{env}}}\rho_{n\tau} = \text{Tr}_{\mathcal{H}_{\text{env}}}U(n\tau, 0)(\rho_p \otimes \rho_\beta^{\otimes M})U(n\tau, 0)^*. \quad (2.15)$$

In fact, for reasons that will become clear later (see Sections 5 and 7), we shall consider the more general linear operator defined on $\mathcal{B}(\mathcal{H}_p)$ by

$$D_{\alpha,n}(A) = \text{Tr}_{\mathcal{H}_{\text{env}}}\left(I \otimes [\rho_\beta^\alpha]^{\otimes M}\right)U(n\tau, 0)\left(A \otimes [\rho_\beta^{1-\alpha}]^{\otimes M}\right)U(n\tau, 0)^*,$$

where α is an arbitrary real parameter. Using the cyclicity of the partial trace w.r.t. atomic operators and the fact that ρ_β is invariant under the free atomic dynamics we can replace $U(n\tau, 0)$ by $e^{-i\tau H_n}U((n-1)\tau, 0)$ in the last formula. It then follows that

$$D_{\alpha,n}(A) = \text{Tr}_{\mathcal{H}_{a,n}}(I \otimes \rho_\beta^\alpha)e^{-i\tau H_n}(D_{\alpha,n-1}(A) \otimes \rho_\beta^{1-\alpha})e^{i\tau H_n} = \mathcal{L}_\alpha(D_{\alpha,n-1}(A)),$$

where

$$\mathcal{L}_\alpha(A) = \text{Tr}_{\mathcal{H}_a}(I \otimes \rho_\beta^\alpha)e^{-i\tau H}(A \otimes \rho_\beta^{1-\alpha})e^{i\tau H}. \quad (2.16)$$

Here we have dropped the index n since $\mathcal{L}_\alpha : \mathcal{B}(\mathcal{H}_p) \rightarrow \mathcal{B}(\mathcal{H}_p)$ does not depend on it. We conclude that $D_{\alpha,n}(A) = \mathcal{L}_\alpha^n(A)$, so that for any B one can write

$$\text{Tr}(B \otimes [\rho_\beta^\alpha]^{\otimes M})U(n\tau, 0)(A \otimes [\rho_\beta^{1-\alpha}]^{\otimes M})U(n\tau, 0)^* = \text{Tr}_{\mathcal{H}_p}B\mathcal{L}_\alpha^n(A). \quad (2.17)$$

The operator \mathcal{L}_α will play a central role in our analysis. For later reference, we describe its main properties in the following proposition.

Proposition 2.1 For any $\alpha \in \mathbb{R}$, \mathcal{L}_α is a completely positive operator on $\mathcal{B}^1(\mathcal{H}_p)$ with spectral radius

$$\theta(\alpha) = (1 - p) + p \frac{\cosh\left(\left(\frac{1}{2} - \alpha\right)\beta E\right)}{\cosh\left(\frac{1}{2}\beta E\right)}, \quad (2.18)$$

where p is defined in (3.2). Its adjoint w.r.t. the duality $\langle A|B \rangle = \text{Tr}AB$ is the completely positive operator on $\mathcal{B}(\mathcal{H}_p)$ given by

$$\mathcal{L}_\alpha^*(B) = \text{Tr}_{\mathcal{H}_a}(I \otimes \rho_\beta^{1-\alpha}) e^{i\tau H} (B \otimes \rho_\beta^\alpha) e^{-i\tau H}.$$

Going back to the special case $\alpha = 0$, we have

$$\rho_{p,n\tau} = \mathcal{L}_0^n(\rho_p),$$

so that the discrete semi-group $(\mathcal{L}_0^n)_{n \in \mathbb{N}}$ acting on the density matrices of \mathcal{H}_p describes the reduced Schrödinger dynamics of the particle. The reduced Heisenberg dynamics is obtained by duality: for $B \in \mathcal{B}(\mathcal{H}_p)$ and $\rho_p \in \mathcal{B}^1(\mathcal{H}_p)$,

$$\text{Tr}_{\mathcal{H}_p} B \mathcal{L}_0^n(\rho_p) = \text{Tr}_{\mathcal{H}_p} \mathcal{L}_0^{*n}(B) \rho_p.$$

At this point, the choice of M becomes immaterial and we can consider an arbitrary large number of interactions. Given an observable B on \mathcal{H}_p , we write

$$\langle B \rangle_n = \text{Tr} B \mathcal{L}_0^n(\rho_p),$$

for its expectation value at time $t = n\tau$.

3 Results and discussion

We are now in a position to state our main results on the dynamics of the particle. As will be shown in Section 4, coupling with a single 2-level atom turns the periodic Bloch oscillations (2.5) of frequency $\omega_{\text{Bloch}} = F$ into quasi-periodic motion with the two frequencies ω_{Bloch} and

$$\omega_0 = \sqrt{(E - F)^2 + 4\lambda^2}, \quad (3.1)$$

(see Equ. (4.3) below). Repeated interactions with 2-level atoms have a much more drastic effect. The bounded motion of the particle now becomes diffusive. In terms of the parameter

$$p = \frac{4\lambda^2}{\omega_0^2} \sin^2\left(\frac{\omega_0\tau}{2}\right) \in [0, 1], \quad (3.2)$$

which is a simple function of the model parameters λ , E and F , the motion is characterized by a drift velocity

$$v_d = v_d(E, F) = \frac{p}{\tau} \tanh\left(\frac{\beta E}{2}\right), \quad (3.3)$$

and a diffusion constant

$$D = D(E, F) = \frac{p}{2\tau} \left(1 - p \tanh^2\left(\frac{\beta E}{2}\right)\right). \quad (3.4)$$

More precisely, the following holds.

Theorem 3.1 *Assume that $F > 0$, $\lambda \neq 0$ and $\omega_0\tau \notin 2\pi\mathbb{Z}$ so that $p \in]0, 1[$. Let the density matrix $\rho_p \in \mathcal{B}^1(\mathcal{H}_p)$ describe the initial state of the particle and denote by μ_n the spectral measures of the position observable X in the state $\rho_{p, n\tau}$,*

$$\mu_n(f) = \int f(x) d\mu_n(x) = \langle f(X) \rangle_n. \quad (3.5)$$

1. *The Central Limit Theorem (CLT) holds: For any bounded continuous f on \mathbb{R} ,*

$$\lim_{n \rightarrow \infty} \int f\left(\frac{x - v_d n\tau}{\sqrt{2Dn\tau}}\right) d\mu_n(x) = \int f(x) e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}.$$

2. *If $\text{Tr}(X^2 \rho_p) < +\infty$, then*

$$\lim_{n \rightarrow \infty} \frac{\langle X \rangle_n}{n\tau} = v_d, \quad \lim_{n \rightarrow \infty} \frac{\langle (X - v_d n\tau)^2 \rangle_n}{n\tau} = 2D.$$

3. *If $\text{Tr}(e^{\gamma|X|} \rho_p) < +\infty$ for all $\gamma > 0$ then a Large Deviation Principle (LDP) holds in the sense that, for any interval $J \subset \mathbb{R}$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mu_n(nJ) = - \inf_{x \in J} I(x), \quad (3.6)$$

where $I(x)$ is the Legendre-Fenchel transform of $e(\eta) = \log \theta(-\eta/\beta E)$, i.e.,

$$I(x) = \sup_{\eta \in \mathbb{R}} \left[\eta x - \log \theta\left(-\frac{\eta}{\beta E}\right) \right].$$

Here the function θ is defined in (2.18).




Figure 1: The cumulant generating function and the large deviation rate function.

Note that when $E = F$, the mobility μ , defined by

$$\mu = \lim_{F \rightarrow 0} \frac{v_d}{F} = \frac{\beta \sin^2(\lambda\tau)}{2\tau},$$

and the diffusion constant

$$D = \mu\beta^{-1} \left(1 - \sin^2(\lambda\tau) \tanh^2 \left(\frac{\beta F}{2} \right) \right),$$

satisfy the Einstein relation

$$\lim_{F \rightarrow 0} D = \mu\beta^{-1} = \mu k_B T.$$

When $E \neq F$, the Einstein relation still holds provided one takes the limit $E \rightarrow 0$ at fixed F (also in the definition of the mobility).

The rate function in Part 3 is explicitly given by

$$I(x) = \begin{cases} -x \left(\frac{\beta E}{2} + \log \left(\frac{R(x) - x}{a(1-x)} \right) \right) - \log \left(\frac{(1-p)(R(x)+1)}{1-x^2} \right) & \text{for } x \in [-1, 1], \\ +\infty & \text{otherwise,} \end{cases}$$

where

$$a = \frac{p}{(1-p) \cosh(\beta E/2)}, \quad R(x) = \sqrt{x^2 + a^2(1-x^2)}.$$

It is strictly convex on $[-1, 1]$ and satisfies $I(v_d\tau) = 0$ and $I(x) > 0$ for $x \neq v_d\tau$ (see Figure 1).

Note that the drift velocity and diffusion constant do not depend on the initial state of the particle. The CLT gives us the probability to find the particle at time $n\tau$ in a region of size $O(\sqrt{n})$ around the mean value $v_d n\tau$, whereas the LDP gives information on

this probability for a region of size $O(n)$. To put it differently, it yields information on the probability that the particle's mean speed falls asymptotically in an interval of size $O(1)$. Loosely speaking, it says that

$$\mu_n(\{n(v_d + \delta v)\tau\}) \simeq e^{-nI((v_d + \delta v)\tau)}.$$

The peculiar symmetry $e(-\beta E - \eta) = e(\eta)$ immediately leads to the relation $I(x) = -\beta E x + I(-x)$ which tells us that

$$\lim_{\delta v \downarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n\tau} \log \frac{\mu_n(n[-v - \delta v, -v + \delta v]\tau)}{\mu_n(n[v - \delta v, v + \delta v]\tau)} = -\beta E v,$$

i.e., that negative mean velocities are exponentially less likely than positive ones. The reader familiar with recent developments in non-equilibrium statistical mechanics will recognize here a kind of fluctuation theorem. Indeed, we shall see that the symmetry of the function $e(\eta)$ is a direct consequence of time-reversal invariance and, in a sense, a remnant of the Evans-Searles (or transient) fluctuation theorem (see (3.7)).

We have further studied the statistics of the energy changes of the particle, the environment and the whole system. Note that the latter is not expected to vanish, since the Hamiltonian is time-dependent, so that total energy is not conserved.

To study the change in the energy of the atomic reservoir we use the following operational procedure. The reservoir being initially in thermal equilibrium at inverse temperature β and the particle in the state ρ_p , we measure the total energy of the reservoir and the particle just before the first interaction and just after the n -th interaction. These successive measurements yield the four values $E_{p,0}, E_{p,n} \in \text{sp } H_p$ and $E_{\text{env},0}, E_{\text{env},n} \in \text{sp } H_{\text{env}}$. It will be convenient to express the resulting change in energy in terms of the ‘‘entropy like’’ quantities

$$\Delta S_{p,n} = \beta^*(E_{p,n} - E_{p,0}), \quad \Delta S_{\text{env},n} = -\beta(E_{\text{env},n} - E_{\text{env},0}),$$

where $\beta^* = \beta E/F$. We denote by \mathbb{P}^n the joint probability distribution of $\Delta S_{p,n}$, $\Delta S_{\text{env},n}$ and by \mathbb{E}^n the corresponding expectation. Note that the quantities $E_{\text{env},0}$ and $E_{\text{env},n}$ are well defined provided the reservoir contains only a finite number M of atoms and $n \leq M$. However, under those circumstances, $\Delta S_{\text{env},n}$ does not depend on M and we can therefore consider the reservoir contains an infinite number of atoms. This simple thermodynamical limit will always be understood in what follows.

Remark. When applied to electric charge, or more generally to particle number, the two measurement processes described above go under the name *full counting statistics* (see e.g. [ABGK] and references therein). The present application is closer to the approach to current fluctuations found in [dR].

Theorem 3.2 1. $\mathbb{P}^n[\Delta S_{p,n} = \Delta S_{\text{env},n}] = 1$. Hence, in the following we set

$$\Delta S_n = \Delta S_{p,n} = \Delta S_{\text{env},n}.$$

2. The cumulant generating function of ΔS_n is given by

$$\log \mathbb{E}^n [e^{\alpha \Delta S_n}] = n \log \theta(\alpha),$$

where the function θ is defined in (2.18).

3. Its mean value and variance are

$$\mathbb{E}^n \left[\frac{\Delta S_n}{n} \right] = -\beta E v_d \tau, \quad \mathbb{E}^n \left[\frac{(\Delta S_n + \beta E v_d n \tau)^2}{n} \right] = (\beta E)^2 2D\tau.$$

4. The CLT holds: For any bounded continuous function f ,

$$\lim_{n \rightarrow \infty} \mathbb{E}^n \left[f \left(\frac{\Delta S_n + \beta E v_d n \tau}{\beta E \sqrt{2Dn\tau}} \right) \right] = \int f(x) e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}.$$

5. The sequence $(\mathbb{P}^n)_{n \in \mathbb{N}}$ satisfies a LDP: For any intervals $J \subset \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{P}^n \left[\frac{\Delta S_n}{n} \in J \right] = - \inf_{s \in J} \phi(s),$$

with the rate function $\phi(s) = \sup_{\alpha \in \mathbb{R}} (\alpha s - \log \theta(\alpha)) = I \left(-\frac{s}{\beta E} \right)$.

6. It satisfies the transient fluctuation theorem

$$\frac{\mathbb{P}^n \left[\frac{\Delta S_n}{n} = -s \right]}{\mathbb{P}^n \left[\frac{\Delta S_n}{n} = s \right]} = e^{ns}. \quad (3.7)$$

Part 1 clearly reflects the fact that the number operator (2.11) commutes with H so that

$$\beta^* H_p + \beta H_{\text{env}} = \beta E \left(\frac{H_p}{F} + \frac{H_{\text{env}}}{E} \right),$$

is preserved by the repeated interaction dynamics.

The particle's drift velocity is v_d , and one sees therefore that, as expected, its energy loss per unit time equals the work done by F per unit time. Simultaneously, the environment gains energy at a rate $E v_d$: indeed, the particle moves on average v_d steps to the right per unit time, which corresponds to v_d elements of the chain gaining an energy E . This

leads to an average energy gain or loss of $(E - F)v_d$ for the full system. In the special case $E = F$, these rates are equal, and the total system neither loses nor gains energy. This is a consequence of the fact (mentioned after Equ. (2.11)) that the interaction term in the Hamiltonian commutes with the free Hamiltonian in this case. In general, the total energy is not preserved, which is a reflection of the fact that the Hamiltonian of the total system is time-dependent, as is clear from (2.13).

Note also that the symmetry $\theta(1 - \alpha) = \theta(\alpha)$ which leads to the transient fluctuation theorem (3.7) is evident from Proposition 2.1. We shall see in Section 5.2 that it is actually a consequence of time-reversal invariance.

Since

$$-\frac{H_p}{F} \simeq X,$$

we expect a very similar result for the position increment $\Delta X_n = X_n - X_0$ obtained from a double measurement of X at time $t = 0$ and $t = n\tau$. Indeed, the distribution \mathbb{Q}^n of ΔX_n satisfies the following.

Theorem 3.3 1. *The cumulant generating function of ΔX_n satisfies*

$$g(\eta) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{Q}^n [e^{\eta \Delta X_n}] = \log \theta \left(-\frac{\eta}{\beta E} \right).$$

2. *Its mean value and variance are*

$$\lim_{n \rightarrow \infty} \mathbb{Q}^n \left[\frac{\Delta X_n}{n\tau} \right] = v_d, \quad \mathbb{Q}^n \left[\frac{(\Delta X_n - v_d n\tau)^2}{n\tau} \right] = 2D.$$

3. *The CLT holds: For any bounded continuous function f ,*

$$\lim_{n \rightarrow \infty} \mathbb{Q}^n \left[f \left(\frac{\Delta X_n - v_d n\tau}{\sqrt{2Dn\tau}} \right) \right] = \int f(x) e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}.$$

4. *The sequence $(\mathbb{Q}^n)_{n \in \mathbb{N}}$ satisfies a LDP: For any intervals $J \subset \mathbb{R}$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{Q}^n \left[\frac{\Delta X_n}{n} \in J \right] = - \inf_{x \in J} I(x).$$

5. *It satisfies the asymptotic fluctuation theorem*

$$\beta E(v - \delta v) \leq \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\mathbb{Q}^n \left[\frac{\Delta X_n}{n\tau} \in [-v - \delta v, -v + \delta v] \right]}{\mathbb{Q}^n \left[\frac{\Delta X_n}{n\tau} \in [v - \delta v, v + \delta v] \right]} \leq \beta E(v + \delta v),$$

for $v \in]-1, 1[$ and $\delta v > 0$, small enough.

Remark. This fluctuation theorem is not of transient (or Evans-Searles) type since it only holds in the large time limit. It is not of the stationary (or Gallavotti-Cohen) type either since there is no stationary state for the dynamics, as we shall see in Section 5.3. For more details on these fluctuation theorems, we refer the reader to e.g. [JPR, RM].

Note also the similarity with Theorem 3.1. Theorem 3.3 describes the position increment of the particle without requiring sharp localization of the particle position at time 0, contrary to Theorem 3.1.

4 Interaction with a single two level atom

In this section we investigate the dynamics of the particle interacting with a single atom described by the Hamiltonian (2.10). As already remarked, H can easily be diagonalized by exploiting the fact that it commutes with the number operator (2.11). To work out this diagonalisation explicitly and to get a tractable formula for the propagator e^{itH} we introduce the unitary operator

$$U = (Tb^*b + bb^*) \cos \theta - (Tb^* - b) \sin \theta,$$

where θ is given by

$$\cos(2\theta) = \frac{E - F}{\omega_0}, \quad \sin(2\theta) = \frac{2\lambda}{\omega_0}.$$

Using the commutation relations (2.2) and (2.6), one then easily checks that

$$U^* H U = H_p + \omega_0 \left(b^* b - \frac{1}{2} \right) + \frac{E - F}{2}. \quad (4.1)$$

It follows immediately that, for $F \neq 0$, the spectrum of H is purely discrete,

$$\text{sp}(H) = 2 - F\mathbb{Z} + \frac{E - F}{2} \pm \frac{\omega_0}{2}.$$

If ω_0/F is not an integer, all the eigenvalues are simple. The normalized eigenvectors are given by

$$\begin{aligned} |\phi_{k,-}\rangle &= U|\psi_k\rangle \otimes |0\rangle = \cos \theta |\psi_k\rangle \otimes |0\rangle - \sin \theta |\psi_{k+1}\rangle \otimes |1\rangle, \\ |\phi_{k,+}\rangle &= U|\psi_k\rangle \otimes |1\rangle = \cos \theta |\psi_{k+1}\rangle \otimes |1\rangle + \sin \theta |\psi_k\rangle \otimes |0\rangle, \end{aligned}$$

where $|0\rangle$ and $|1\rangle$ denote the ground state and the excited state of the atom. Of course the situation is completely different when $F = 0$. The spectrum

$$\text{sp}(H) = \left(\frac{1}{2}(E - F - \omega_0) + [0, 4] \right) \cup \left(\frac{1}{2}(E - F + \omega_0) + [0, 4] \right),$$

is then purely absolutely continuous.

From Equ. (4.1) we get the explicit formula for the propagator,

$$e^{itH} = U e^{it(E-F)/2} e^{it\omega_0(b^*b-1/2)} e^{itH_p} U^*, \quad (4.2)$$

which reduces the proof of the following result to a straightforward calculation.

Lemma 4.1 *For any operator A on \mathcal{H}_p and any $t \in \mathbb{R}$ one has*

$$e^{-itH} A \otimes \rho_\beta e^{itH} = \mathcal{A}_{\beta,t}(A_t) \otimes b^*b + \mathcal{B}_{\beta,t}(A_t) \otimes b + \mathcal{B}_{\beta,t}(A_t^*)^* \otimes b^* + \mathcal{C}_{\beta,t}(A_t) \otimes bb^*,$$

where $A_t = e^{-itH_p} A e^{itH_p}$ and

$$\begin{aligned} \mathcal{A}_{\beta,t}(A) &= \frac{e^{-\beta E}}{Z_\beta} \left(1 - \frac{4\lambda^2}{\omega_0^2} \sin^2 \left(\frac{\omega_0 t}{2} \right) \right) A + \frac{1}{Z_\beta} \frac{4\lambda^2}{\omega_0^2} \sin^2 \left(\frac{\omega_0 t}{2} \right) T A T^*, \\ \mathcal{B}_{\beta,t}(A) &= \frac{1}{Z_\beta} \frac{2\lambda}{\omega_0} \left(\frac{i}{2} \sin(\omega_0 t) - \frac{E-F}{\omega_0} \sin^2 \left(\frac{\omega_0 t}{2} \right) \right) (A T^* - e^{-\beta E} T^* A), \\ \mathcal{C}_{\beta,t}(A) &= \frac{1}{Z_\beta} \left(1 - \frac{4\lambda^2}{\omega_0^2} \sin^2 \left(\frac{\omega_0 t}{2} \right) \right) A + \frac{e^{-\beta E}}{Z_\beta} \frac{4\lambda^2}{\omega_0^2} \sin^2 \left(\frac{\omega_0 t}{2} \right) T^* A T. \end{aligned}$$

By noting that $X = 2X \otimes \rho_{\beta=0}$ the preceding lemma can be used to compute the evolution of the position observable

$$\begin{aligned} e^{itH} X e^{-itH} &= e^{itH_p} X e^{-itH_p} \\ &+ \left(\frac{4\lambda^2}{\omega_0^2} (bb^* - b^*b) + \frac{2\lambda(E-F)}{\omega_0^2} (Tb^* + T^*b) \right) \sin^2 \left(\frac{\omega_0 t}{2} \right) \\ &- i \frac{\lambda}{\omega_0} (Tb^* - T^*b) \sin(\omega_0 t). \end{aligned} \quad (4.3)$$

We conclude that the coupling to a single atom does not substantially alter the long term behavior of the particle. In particular, when $F \neq 0$, the motion remains bounded. We will see in the next section that the situation is very different for repeated interactions with a sequence of atoms.

5 Repeated interaction dynamics

In this section we study the properties of the operator \mathcal{L}_α , defined in (2.16). Besides the special case $\alpha = 0$ which describes the reduced Schrödinger dynamics of the particle and is of course central, the operator \mathcal{L}_α appears in the study of the cumulant generating

function of the “entropy” (see Theorem 3.2 and (7.3) in Section 7.1). In particular we shall see that the consequence of the time reversal invariance of the dynamics of the particle coupled to a two-level atom on \mathcal{L}_α is at the origin of the symmetry $\theta(1 - \alpha) = \theta(\alpha)$ and hence of the transient fluctuation theorem (3.7).

It is clear that $\alpha \mapsto \mathcal{L}_\alpha$ is entire analytic as a map from \mathbb{C} to the bounded operators on $\mathcal{B}(\mathcal{H}_p)$. Moreover, since

$$\mathrm{Tr}_{\mathcal{H}_p} B \mathcal{L}_\alpha(A) = \mathrm{Tr}(B \otimes \rho_\beta^\alpha) e^{-i\tau H} (A \otimes \rho_\beta^{1-\alpha}) e^{i\tau H},$$

\mathcal{L}_α is also bounded as an operator on the Schatten-von Neumann class $\mathcal{B}^p(\mathcal{H}_p)$, for any $p \in [1, \infty]$.

Using the cyclicity of the partial trace w.r.t. operators on \mathcal{H}_a we can write, for $\alpha \in \mathbb{R}$,

$$\begin{aligned} \mathcal{L}_\alpha(A) &= \mathrm{Tr}_{\mathcal{H}_a} \left((I \otimes \rho_\beta^\alpha) e^{-i\tau H} (A \otimes \rho_\beta^{1-\alpha}) e^{i\tau H} \right) \\ &= \mathrm{Tr}_{\mathcal{H}_a} \left((I \otimes e^{-\alpha\beta H_a/2}) e^{-i\tau H} (I \otimes e^{\alpha\beta H_a/2}) \times (A \otimes \rho_\beta) \right. \\ &\quad \left. \times (I \otimes e^{\alpha\beta H_a/2}) e^{i\tau H} (I \otimes e^{-\alpha\beta H_a/2}) \right) \\ &= \mathrm{Tr}_{\mathcal{H}_a} \left(e^{-i\tau H^{(\alpha)}} A \otimes \rho_\beta e^{i\tau H^{(\alpha)*}} \right), \end{aligned} \quad (5.1)$$

where $H^{(\alpha)}$ is a bounded perturbation of H ,

$$\begin{aligned} H^{(\alpha)} &= e^{-\alpha\beta H_a/2} H e^{\alpha\beta H_a/2} \\ &= H_p + H_a + \lambda e^{-\alpha\beta H_a/2} V e^{\alpha\beta H_a/2} \\ &= H_p + H_a + \lambda (e^{-\alpha\beta E/2} T b^* + e^{\alpha\beta E/2} T^* b). \end{aligned} \quad (5.2)$$

This shows in particular that \mathcal{L}_α is completely positive for real values of α .

5.1 Gauge invariance

Lemma 5.1 *The operator \mathcal{L}_α commutes with the evolution of the non-interacting particle, i.e.,*

$$\mathcal{L}_\alpha \left(e^{-itH_p} A e^{itH_p} \right) = e^{-itH_p} \mathcal{L}_\alpha(A) e^{itH_p}, \quad (5.3)$$

holds for all $t, \alpha \in \mathbb{R}$ and $A \in \mathcal{B}(\mathcal{H}_p)$.

Proof. From the fact that H commutes with the number operator (2.11) we infer

$$H = e^{itN} H e^{-itN} = e^{itH_p/F} e^{itH_a/E} H e^{-itH_a/E} e^{-itH_p/F},$$

so that

$$e^{-itH_a/E} e^{-i\tau H} e^{itH_a/E} = e^{itH_p/F} e^{-i\tau H} e^{-itH_p/F}.$$

Since H_p and H_a commute, we also have

$$e^{-itH_a/E} e^{-i\tau H^{(\alpha)}} e^{itH_a/E} = e^{itH_p/F} e^{-i\tau H^{(\alpha)}} e^{-itH_p/F}, \quad (5.4)$$

where $H^{(\alpha)}$ is defined in (5.2), and hence

$$\begin{aligned} & e^{itH_p} \mathcal{L}_\alpha \left(e^{-itH_p} A e^{itH_p} \right) e^{-itH_p} \\ &= \text{Tr}_{\mathcal{H}_a} \left(e^{itH_p} e^{-i\tau H^{(\alpha)}} e^{-itH_p} A \otimes \rho_\beta e^{itH_p} e^{i\tau H^{(\alpha)*}} e^{-itH_p} \right) \\ &= \text{Tr}_{\mathcal{H}_a} \left(e^{-itH_a F/E} e^{-i\tau H^{(\alpha)}} e^{itH_a F/E} A \otimes \rho_\beta e^{-itH_a F/E} e^{i\tau H^{(\alpha)*}} e^{itH_a F/E} \right) \\ &= \text{Tr}_{\mathcal{H}_a} \left(e^{-itH_a F/E} e^{-i\tau H^{(\alpha)}} A \otimes \rho_\beta e^{i\tau H^{(\alpha)*}} e^{itH_a F/E} \right) \\ &= \mathcal{L}_\alpha(A), \end{aligned}$$

where we used (5.1) in the 1st line, (5.4) in the 2nd one, the fact that H_a commutes with $A \otimes \rho_\beta$ in the 3rd one, and finally the cyclicity of the trace. \square

Introducing the non-interacting evolution operator

$$\mathcal{U}(A) = e^{-i\tau H_p} A e^{i\tau H_p},$$

we define the ‘‘interaction picture’’ reduced evolution as $\tilde{\mathcal{L}}_\alpha = \mathcal{L}_\alpha \circ \mathcal{U}^{-1}$. Note that, by Lemma 5.1, we have

$$\mathcal{L}_\alpha^n = \tilde{\mathcal{L}}_\alpha^n \circ \mathcal{U}^n = \mathcal{U}^n \circ \tilde{\mathcal{L}}_\alpha^n,$$

for any $n \in \mathbb{N}$.

5.2 Time-reversal invariance

Let us denote by C_p the complex conjugation on $\ell^2(\mathbb{Z})$, i.e., $(C_p \psi)(x) = \overline{\psi(x)}$ and set

$$\mathcal{C}(A) = C_p A C_p^*,$$

for $A \in \mathcal{B}(\mathcal{H}_p)$. This anti-linear involution implements time reversal of the particle’s dynamics. Indeed, since the Hamiltonian H_p is real in the position representation, $C_p H_p = H_p C_p$, one has $C_p e^{itH_p} = e^{-itH_p} C_p$ for all $t \in \mathbb{R}$ and in particular

$$\mathcal{C} \circ \mathcal{U} = \mathcal{U}^{-1} \circ \mathcal{C}.$$

Lemma 5.2 *For all $\alpha \in \mathbb{R}$ one has $\mathcal{L}_\alpha^* = \mathcal{C} \circ \mathcal{L}_{1-\alpha} \circ \mathcal{C}$.*

Remark. As will be clear from its proof, this property of \mathcal{L}_α is a consequence of the time-reversal invariance of the dynamics of the particle coupled to a two-level atom. It implies that the spectral radius of \mathcal{L}_α and $\mathcal{L}_{1-\alpha}$ coincide.

Proof. Setting $C_a(a_0|0\rangle + a_1|1\rangle) = \overline{a_0}|0\rangle + \overline{a_1}|1\rangle$ defines a anti-unitary operator on \mathcal{H}_a such that $C_a H_a = H_a C_a$ and hence

$$C_a e^{itH_a} = e^{-itH_a} C_a, \quad C_a \rho_\beta = \rho_\beta C_a.$$

Since T and T^* are real w.r.t. C_p and b and b^* are real w.r.t. C_a , the Hamiltonian H is real w.r.t. $C = C_p \otimes C_a$ and one has $C e^{itH} = e^{-itH} C$.

From the fact that the partial trace on \mathcal{H}_a satisfies

$$\text{Tr}_{\mathcal{H}_a} C A C^* = \mathcal{C}(\text{Tr}_{\mathcal{H}_a} A),$$

for all $A \in \mathcal{B}(\mathcal{H}_p \otimes \mathcal{H}_a)$, one deduces that, for $A, B \in \mathcal{B}(\mathcal{H}_p)$,

$$\begin{aligned} \mathcal{C}(A \mathcal{L}_\alpha(B)) &= \mathcal{C}(\text{Tr}_{\mathcal{H}_a}(A \otimes \rho_\beta^\alpha) e^{-i\tau H} (B \otimes \rho_\beta^{1-\alpha}) e^{i\tau H}) \\ &= \text{Tr}_{\mathcal{H}_a} C(A \otimes \rho_\beta^\alpha) e^{-i\tau H} (B \otimes \rho_\beta^{1-\alpha}) e^{i\tau H} C^* \\ &= \text{Tr}_{\mathcal{H}_a} (\mathcal{C}(A) \otimes \rho_\beta^\alpha) e^{i\tau H} (\mathcal{C}(B) \otimes \rho_\beta^{1-\alpha}) e^{-i\tau H} \\ &= \text{Tr}_{\mathcal{H}_a} (\mathcal{C}(B) \otimes \rho_\beta^{1-\alpha}) e^{-i\tau H} (\mathcal{C}(A) \otimes \rho_\beta^\alpha) e^{i\tau H} \\ &= \mathcal{C}(B) \mathcal{L}_{1-\alpha} \circ \mathcal{C}(A). \end{aligned}$$

It follows that

$$\begin{aligned} \text{Tr}_{\mathcal{H}_p} A \mathcal{L}_\alpha(B) &= \overline{\text{Tr}_{\mathcal{H}_p} \mathcal{C}(A \mathcal{L}_\alpha(B))} \\ &= \overline{\text{Tr}_{\mathcal{H}_p} \mathcal{C}(B) \mathcal{L}_{1-\alpha} \circ \mathcal{C}(A)} \\ &= \text{Tr}_{\mathcal{H}_p} \mathcal{C} \circ \mathcal{L}_{1-\alpha} \circ \mathcal{C}(A) B, \end{aligned}$$

and hence $\mathcal{L}_\alpha^*(A) = \mathcal{C} \circ \mathcal{L}_{1-\alpha} \circ \mathcal{C}(A)$. \square

5.3 Kraus representation

Since \mathcal{L}_α is completely positive for $\alpha \in \mathbb{R}$, the same is true of $\tilde{\mathcal{L}}_\alpha$. The following result describes the Kraus representation of the latter operator.

Lemma 5.3 *For any $\alpha \in \mathbb{R}$ and $A \in \mathcal{B}(\mathcal{H}_p)$, one has*

$$\tilde{\mathcal{L}}_\alpha(A) = e^{\alpha\beta E} p_- T^* A T + p_0 A + e^{-\alpha\beta E} p_+ T A T^*, \quad (5.5)$$

where

$$p_- = \frac{e^{-\beta E}}{1 + e^{-\beta E}} p, \quad p_0 = 1 - p, \quad p_+ = \frac{1}{1 + e^{-\beta E}} p.$$

with p defined in Equ. (3.2).

We note that $e^{\alpha\beta E}p_- + p_0 + e^{-\alpha\beta E}p_+ = \theta(\alpha)$, where $\theta(\alpha)$ is defined in Proposition 2.1, so

$$\mathcal{L}_\alpha(I) = \mathcal{L}_\alpha^*(I) = \theta(\alpha)I. \quad (5.6)$$

Consider $\mathcal{K}_\alpha = \theta(\alpha)^{-1}\mathcal{L}_\alpha$ as an operator on $\mathcal{B}^1(\mathcal{H}_p)$. Since \mathcal{K}_α is completely positive and trace preserving, it has unit spectral radius (see [Sch]). Hence, $\theta(\alpha)$ is the spectral radius of \mathcal{L}_α . This proves Proposition 2.1. The remark after Lemma 5.2 now explains the origin of the symmetry $\theta(1 - \alpha) = \theta(\alpha)$.

Let us now set $\alpha = 0$ in Equ. (5.5) and explore the implications of this expression for the dynamics of the particle. If ρ describes the state of the particle, then $T^*\rho T$ (respectively $T\rho T^*$) represent the same state translated by one lattice spacing to the left (respectively right). Note moreover that

$$p_- + p_0 + p_+ = 1,$$

so that the reduced evolution $\mathcal{L}_0 = \tilde{\mathcal{L}}_0 \circ \mathcal{U}$ consists of a free evolution with the Hamiltonian H_p , followed by a random translation by ± 1 or 0 , and with probabilities p_\pm or p_0 . Note that the dynamics is trivial if $p = 0$, i.e., if $\omega_0\tau = 2\pi m$ with $m \in \mathbb{Z}$. In that case there is no translation and the particle evolves according to H_p . This can be seen directly on Equ. (4.2) by noticing that $UH_pU^* = H_p + Fb^*b$. It follows that the propagator factorizes

$$e^{i\tau H} = (-1)^m e^{i\tau(E-F)/2} e^{i\tau H_p} \otimes e^{i\tau Fb^*b},$$

and, up to an inessential phase factor and a renormalization of the atomic Bohr frequency, the particle and the two-level system evolve as if they were not coupled. This resembles the ‘‘Rabi oscillation’’ phenomenon which appears in the Jaynes-Cummings model for matter-radiation interaction. In the following we will avoid this resonance and assume $p \neq 0$.

Introducing a family of i.i.d. random variables Y_n , taking the values ± 1 and 0 with probability p_\pm and p_0 , and defining $S_n = \sum_{j=1}^n Y_j$, we can very concisely write

$$\mathcal{L}_0^n(\rho) = e^{-in\tau H_p} \mathbb{E} [T^{S_n} \rho T^{-S_n}] e^{in\tau H_p}. \quad (5.7)$$

Accordingly, the study of the dynamics of the system is reduced to that of a classical random walk. As a further remark, suppose that the initial state of the particle is invariant under the uncoupled dynamics, so that

$$\rho = \sum_{k \in \mathbb{Z}} p_k |\psi_k\rangle \langle \psi_k|. \quad (5.8)$$

Then, using Equ. (2.9), we obtain $\mathcal{L}_0^n(\rho) = \sum_{k \in \mathbb{Z}} p_k^{(n)} |\psi_k\rangle \langle \psi_k|$ with

$$p_k^{(n+1)} = p_+ p_{k-1}^{(n)} + p_0 p_k^{(n)} + p_- p_{k+1}^{(n)}. \quad (5.9)$$

Thus, the set of H_p -invariant states is invariant under the reduced dynamics and the latter reduces to a classical Markov chain on this set.

Before turning to the proof of Lemma 5.3, let us show that the reduced dynamics has no stationary state: there exists no density matrix ρ on \mathcal{H}_p such that $\mathcal{L}_0(\rho) = \rho$. Indeed, it follows from Lemma 5.1 that the subspaces \mathcal{J}_d , $d \in \mathbb{Z}$, defined by

$$\begin{aligned}\mathcal{J}_d &= \{\rho \in \mathcal{B}^1(\mathcal{H}_p) \mid e^{-itH_p} \rho e^{itH_p} = e^{itd} \rho \text{ for all } t \in \mathbb{R}\} \\ &= \{\rho \in \mathcal{B}^1(\mathcal{H}_p) \mid \rho = \sum_{k \in \mathbb{Z}} \rho_k |\psi_k\rangle \langle \psi_{k+d}|\},\end{aligned}$$

are globally invariant under \mathcal{L}_0 . Hence, if a state ρ is stationary, so is its diagonal part $\rho_0 = \sum_k p_k |\psi_k\rangle \langle \psi_k|$, where $p_k = \langle \psi_k | \rho | \psi_k \rangle$. Equ. (5.9) then writes

$$p_{k-1} - Z_\beta p_k + e^{-\beta E} p_{k+1} = 0,$$

which implies that $p_k = a + b e^{\beta E k}$ for some constants $a, b \in \mathbb{R}$. But this contradicts the fact that $1 = \text{Tr } \rho = \sum_k p_k$.

Proof of Lemma 5.3. We start with the fact that

$$\rho_\beta^\alpha e^{-i\tau H} A \otimes \rho_\beta^{1-\alpha} e^{i\tau H} = \frac{Z_{\alpha\beta} Z_{(1-\alpha)\beta}}{Z_\beta} \rho_{\alpha\beta} e^{-i\tau H} A \otimes \rho_{(1-\alpha)\beta} e^{i\tau H},$$

so that, applying Lemma 4.1, we get

$$\begin{aligned}\rho_\beta^\alpha e^{-i\tau H} A \otimes \rho_\beta^{1-\alpha} e^{i\tau H} &= \frac{Z_{\alpha\beta} Z_{(1-\alpha)\beta}}{Z_\beta} \rho_{\alpha\beta} (\mathcal{A}_{(1-\alpha)\beta,t}(A_t) b^* b + \mathcal{B}_{(1-\alpha)\beta,t}(A_t) b \\ &\quad + \mathcal{B}_{(1-\alpha)\beta,t}(A_t)^* b^* + \mathcal{C}_{(1-\alpha)\beta,t}(A_t) b b^*).\end{aligned}$$

Upon taking the partial trace over \mathcal{H}_a , we obtain

$$\mathcal{L}_\alpha(A) = \frac{Z_{\alpha\beta} Z_{(1-\alpha)\beta}}{Z_\beta} \left(\mathcal{A}_{(1-\alpha)\beta,t}(A_t) \frac{e^{-\alpha\beta E}}{Z_{\alpha\beta}} + \mathcal{C}_{(1-\alpha)\beta,t}(A_t) \frac{1}{Z_{\alpha\beta}} \right),$$

and the result follows from Lemma 4.1 with some elementary algebra. \square

6 Proof of Theorem 3.1

To complete the proof of our main results, we shall need the following technical lemma.

Lemma 6.1 1. If $F \neq 0$ then, for any $\eta \in \mathbb{R}$, one has

$$\lim_{t \rightarrow \infty} \|e^{itH_p} e^{i\eta X/\sqrt{t}} e^{-itH_p} - e^{i\eta X/\sqrt{t}}\| = 0.$$

2. If $F \neq 0$ then, for any $\eta \in \mathbb{R}$, there exists $C_\eta \geq 1$ such that

$$C_\eta^{-1} \leq e^{-\eta X/2} e^{itH_p} e^{\eta X} e^{-itH_p} e^{-\eta X/2} \leq C_\eta,$$

for all $t \in \mathbb{R}$.

Proof. 1. By Equ. (2.5), one has $e^{itH_p} X e^{-itH_p} = X + B_t$, where B_t is a uniformly bounded operator valued function of t . Duhamel formula yields

$$e^{itH_p} e^{i\eta X/\sqrt{t}} e^{-itH_p} - e^{i\eta X/\sqrt{t}} = \frac{\eta}{\sqrt{t}} R_t,$$

where

$$R_t = \int_0^1 e^{i(1-s)\eta(X+B_t)/\sqrt{t}} B_t e^{is\eta X/\sqrt{t}} ds.$$

The claim follows from the fact that R_t is also uniformly bounded.

2. By Equ. (2.3) we can write, for any $\eta, t \in \mathbb{R}$,

$$Q(t, \eta) = e^{itH_p} e^{-i\eta X} e^{-itH_p} e^{i\eta X} = e^{iG} e^{-itFX} e^{-iG} e^{-i\eta X} e^{iG} e^{itFX} e^{-iG} e^{i\eta X}.$$

From the commutation relation (2.2) we get, for $\theta \in \mathbb{R}$,

$$e^{-i\theta X} G e^{i\theta X} = \frac{2}{F} \sin(\xi + \theta),$$

so that

$$Q(t, \eta) = e^{2i(\sin \xi - \sin(\xi+tF) - \sin(\xi+\eta) + \sin(\xi+tF+\eta))/F}.$$

It follows that $\eta \mapsto Q(t, \eta) \in \mathcal{B}(\mathcal{H}_p)$ extends to an entire analytic function. Moreover, one easily shows that

$$C_\eta = \sup_{t \in \mathbb{R}} \|Q(t, \eta)\| = e^{4 \sinh(|\operatorname{Im} \eta|)/F},$$

for any $\eta \in \mathbb{C}$.

Now since $X_t = e^{itH_p} X e^{-itH_p}$ is self-adjoint the subspace $\mathcal{D}_t = \operatorname{Dom}(e^{X_t^2})$ is dense in \mathcal{H}_p and such that, for $\phi \in \mathcal{D}_t$, the vector valued function $\eta \mapsto e^{i\eta X_t} \phi$ is entire analytic. For $\psi \in \mathcal{D}_t$, $\phi \in \mathcal{D}_0$ and $\eta \in \mathbb{R}$ one has

$$\langle e^{i\bar{\eta} X_t} \psi | e^{i\eta X} \phi \rangle = \langle \psi | Q(t, \eta) \phi \rangle.$$

By analytic continuation, both sides of this identity extend to complex values of η . In particular, one has

$$\langle e^{\eta X_t} \psi | e^{-\eta X} \phi \rangle = \langle \psi | Q(t, i\eta) \phi \rangle,$$

for any $\eta \in \mathbb{R}$. Since \mathcal{D}_t is a core of $e^{\pm \eta X_t}$, the last identity extends to all $\phi \in \text{Dom}(e^{-\eta X})$ and $\psi \in \text{Dom}(e^{\eta X_t})$. The modulus of its right hand side being bounded by $C_{i\eta} \|\psi\| \|\phi\|$, one concludes that $\text{Ran}(e^{-\eta X}) \subset \text{Dom}(e^{\eta X_t})$ and

$$\|e^{\eta X_t} e^{-\eta X}\| \leq C_{i\eta}.$$

It follows that

$$e^{-\eta X/2} e^{\eta X_t} e^{-\eta X/2} = (e^{\eta X_t/2} e^{-\eta X/2})^* (e^{\eta X_t/2} e^{-\eta X/2}) \leq C_{i\eta/2}^2,$$

and

$$(e^{-\eta X/2} e^{\eta X_t} e^{-\eta X/2})^{-1} = e^{\eta X/2} e^{-\eta X_t} e^{\eta X/2} \leq C_{-i\eta/2}^2 = C_{i\eta/2}^2,$$

which together imply

$$C_{i\eta/2}^{-2} \leq e^{-\eta X/2} e^{\eta X_t} e^{-\eta X/2} \leq C_{i\eta/2}^2.$$

□

We now turn to the proof of Theorem 3.1. First, from Equ. (5.7) we find that

$$\langle f(X) \rangle_n = \text{Tr}(f(X) \mathcal{L}_0^n(\rho_p)) = \mathbb{E} [\text{Tr}(f(T^{-S_n} e^{in\tau H_p} X e^{-in\tau H_p} T^{S_n}) \rho_p)].$$

Hence, using Equ. (2.5), we get

$$\begin{aligned} \langle f(X) \rangle_n &= \mathbb{E} [\text{Tr}(f(T^{-S_n} (X + B_n) T^{S_n}) \rho_p)] \\ &= \mathbb{E} [\text{Tr}(f(X + S_n + B_n) \rho_p)], \end{aligned} \tag{6.1}$$

where

$$B_n = \frac{4}{F} \sin\left(\frac{n\tau F}{2}\right) \sin\left(\xi + \frac{n\tau F}{2}\right). \tag{6.2}$$

To prove Part 1, we study the convergence in distribution of the sequence of probability measures $\tilde{\mu}_n(J) = \mu_n(\sqrt{2Dn\tau}J + v_d n\tau)$, $J \subset \mathbb{R}$, where μ_n is defined in (3.5). By the Lévy-Cramér continuity theorem, this is equivalent to the pointwise convergence of the sequence of their characteristic functions. We shall therefore prove that for any $\eta \in \mathbb{R}$

$$\tilde{\mu}_n(e^{i\eta x}) = \int_{\mathbb{R}} e^{i\eta x} d\tilde{\mu}_n(x) = \left\langle e^{i\eta(X - v_d n\tau)/\sqrt{2Dn\tau}} \right\rangle_n \xrightarrow{n \rightarrow \infty} e^{-\eta^2/2}.$$

Using (6.1), we can write

$$\begin{aligned}\tilde{\mu}_n(e^{i\eta x}) &= \mathbb{E} \left[\text{Tr} \left(e^{i\eta(X+S_n-v_d n\tau+B_n)/\sqrt{2Dn\tau}} \rho_p \right) \right] \\ &= \mathbb{E} \left[e^{i\eta(S_n-v_d n\tau)/\sqrt{2Dn\tau}} \right] \text{Tr} \left(e^{i\eta(X+B_n)/\sqrt{2Dn\tau}} \rho_p \right).\end{aligned}\quad (6.3)$$

The classical CLT implies that the first factor of the last line converges, as $n \rightarrow \infty$, to $e^{-\eta^2/2}$ since, as a simple computation confirms

$$\frac{1}{n\tau} \mathbb{E}[S_n] = v_d, \quad \frac{1}{n\tau} \text{Var}[S_n] = 2D, \quad (6.4)$$

where v_d and D are defined in (3.3) and (3.4). The second factor is controlled by Part 1 of Lemma 6.1 which implies

$$\lim_{n \rightarrow \infty} \text{Tr} \left(e^{i\eta(X+B_n)/\sqrt{2Dn\tau}} \rho_p \right) = \lim_{n \rightarrow \infty} \text{Tr} \left(e^{i\eta X/\sqrt{2Dn\tau}} \rho_p \right).$$

The right hand side of this identity is 1 by the dominated convergence theorem.

The first assertion in Part 2 follows from

$$\begin{aligned}\frac{1}{n\tau} \langle X \rangle_n &= \frac{1}{n\tau} \text{Tr}(X \mathcal{L}_0^n(\rho_p)) \\ &= \frac{1}{n\tau} \mathbb{E} [\text{Tr}((X + S_n + B_n)\rho_p)] \\ &= \frac{1}{n\tau} \mathbb{E}[S_n] + \frac{1}{n\tau} \text{Tr}((X + B_n)\rho_p),\end{aligned}$$

since the first term on the last line converges to v_d (by Equ. (6.4)) while the second vanishes as $n \rightarrow \infty$. The proof of the second assertion is similar.

To prove Part 3, we shall study the cumulant generating function

$$e_n(\eta) = \log \mu_n(e^{\eta x}) = \log \text{Tr} \left(e^{\eta X} \mathcal{L}_0^n(\rho_p) \right). \quad (6.5)$$

By Equ. (6.1)-(6.2) and (2.5), one has

$$e_n(\eta) = \log \mathbb{E} \left[\text{Tr} \left(e^{\eta(X+S_n+B_n)} \rho_p \right) \right] = \log \mathbb{E} \left[e^{\eta S_n} \right] + \log \text{Tr} \left(e^{i n \tau H_p} e^{\eta X} e^{-i n \tau H_p} \rho_p \right).$$

One easily computes the first term on the right hand side

$$\log \mathbb{E} \left[e^{\eta S_n} \right] = n \log \mathbb{E} \left[e^{\eta Y_1} \right] = n \log \theta \left(-\frac{\eta}{\beta E} \right).$$

Writing the second term as

$$\log \operatorname{Tr} \left((\rho_{\mathbb{P}}^{1/2} e^{\eta X/2}) e^{-\eta X/2} e^{i n \tau H_{\mathbb{P}}} e^{\eta X} e^{-i n \tau H_{\mathbb{P}}} e^{-\eta X/2} (e^{\eta X/2} \rho_{\mathbb{P}}^{1/2}) \right),$$

and applying Part 2 of Lemma 6.1 we get the bound

$$|\log \operatorname{Tr} (e^{i n \tau H_{\mathbb{P}}} e^{\eta X} e^{-i n \tau H_{\mathbb{P}}} \rho_{\mathbb{P}}) | \leq \log C_{\eta} + \log \operatorname{Tr} (e^{\eta X} \rho_{\mathbb{P}}),$$

from which we conclude that

$$e(\eta) = \lim_{n \rightarrow \infty} \frac{1}{n} e_n(\eta) = \log \theta \left(-\frac{\eta}{\beta E} \right).$$

Since $e(\eta)$ is differentiable, the Gärtner-Ellis theorem (see e.g. [DZ]) implies the LDP (3.6) with the rate function $I(x)$ related to $e(\eta)$ via the Legendre-Fenchel transform. Finally, the symmetry $\theta(1 - \alpha) = \theta(\alpha)$ translates into

$$e(\eta) = e(-\beta E - \eta),$$

which implies $I(x) = -\beta E x + I(-x)$.

7 Full counting statistics

In this section we start with a precise description of the two measurements processes involved in the formulation of Theorems 3.2 and 3.3 and proceed then to the proofs of these results.

Suppose that the initial state of the particle is described by the density matrix $\rho_{\mathbb{P}}$ and set $\rho = \rho_{\mathbb{P}} \otimes \rho_{\beta}^{\otimes M}$. Let A_1, \dots, A_m be commuting self-adjoint operators on the Hilbert space $\mathcal{H} = \mathcal{H}_{\mathbb{P}} \otimes \mathcal{H}_{\text{env}}$. We assume the A_j to have pure point spectrum. We define the vector valued observable $A = (A_1, \dots, A_m)$ and its spectrum $\operatorname{sp} A = \operatorname{sp} A_1 \times \dots \times \operatorname{sp} A_m$. We denote by P_a the spectral projection associated to the eigenvalue $a \in \operatorname{sp} A$.

The outcome of a first measurement of the observables A at time $t = 0$ will be $a \in \operatorname{sp} A$ with probability $\operatorname{Tr} \rho P_a$. After this measurement the state of the combined system is reduced to $\rho' = P_a \rho P_a / \operatorname{Tr}(\rho P_a)$. This state now evolves under the repeated interaction dynamics and, after the n -th interaction, becomes $U(n\tau, 0) \rho' U(n\tau, 0)^*$. A second measurement of A at time $t = n\tau$ will yield the result a' with probability $\operatorname{Tr} U(n\tau, 0) \rho' U(n\tau, 0)^* P_{a'}$. Thus, the joint probability distribution for the two successive measurements of A is given by

$$\begin{aligned} \mathbb{P}_A^n(a, a') &= \operatorname{Tr} \rho P_a \times \operatorname{Tr} U(n\tau, 0) \rho' U(n\tau, 0)^* P_{a'} \\ &= \operatorname{Tr} U(n\tau, 0) P_a \rho P_a U(n\tau, 0)^* P_{a'}. \end{aligned}$$

Therefore, the probability distribution of the measured increment $\Delta a = a' - a$ after n interactions is

$$\mathbb{P}_A^n(\Delta a) = \sum_{\substack{a, a' \in \text{sp } A \\ a' - a = \Delta a}} \text{Tr } U(n\tau, 0) P_a \rho P_a U(n\tau, 0)^* P_{a'}.$$

The cumulant generating function of this distribution is

$$\begin{aligned} g_n(\alpha) &= \log \sum_{\Delta a \in \text{sp } A - \text{sp } A} \mathbb{P}_A^n(\Delta a) e^{\alpha \cdot \Delta a} \\ &= \log \sum_{a, a' \in \text{sp } A} e^{\alpha \cdot (a' - a)} \text{Tr } U(n\tau, 0) P_a \rho P_a U(n\tau, 0)^* P_{a'}, \end{aligned}$$

where \cdot denotes the Euclidean scalar product on \mathbb{R}^m . At this point, it is useful to note that

$$\tilde{\rho} = \sum_{a \in \text{sp } A} P_a \rho P_a,$$

is a density matrix which commutes with A so that $P_a \rho P_a = \tilde{\rho} P_a$. Hence, we can rewrite

$$\begin{aligned} g_n(\alpha) &= \log \sum_{a, a' \in \text{sp } A} e^{\alpha \cdot (a' - a)} \text{Tr } U(n\tau, 0) \tilde{\rho} P_a U(n\tau, 0)^* P_{a'} \\ &= \log \text{Tr } U(n\tau, 0) \tilde{\rho} e^{-\alpha \cdot A} U(n\tau, 0)^* e^{\alpha \cdot A}. \end{aligned} \quad (7.1)$$

In the special case where the A_j are observables of the particle, we obtain

$$g_n(\alpha) = \log \text{Tr } \mathcal{L}_0^n(\tilde{\rho}_p e^{-\alpha \cdot A}) e^{\alpha \cdot A} = \log \text{Tr } \tilde{\rho}_p e^{-\alpha \cdot A} \mathcal{L}_0^{*n}(e^{\alpha \cdot A}), \quad (7.2)$$

where $\tilde{\rho}_p = \sum_{a \in \text{sp } A} P_a \rho_p P_a$.

7.1 Proof of Theorem 3.2

To prove theorem 3.2 we consider the case $A = (\beta^* H_p, -\beta H_{\text{env}})$. From Equ. (7.1), and for any $n \leq M$ (where M is the number of atoms in the reservoir), we have

$$\begin{aligned} g_n(\alpha_p, \alpha_{\text{env}}) &= \log \mathbb{E}^n [e^{\alpha_p \Delta S_{p,n} + \alpha_{\text{env}} \Delta S_{\text{env},n}}] \\ &= \log \text{Tr } U(n\tau, 0) \tilde{\rho} e^{-\alpha_p \beta^* H_p + \alpha_{\text{env}} \beta H_{\text{env}}} U(n\tau, 0)^* e^{\alpha_p \beta^* H_p - \alpha_{\text{env}} \beta H_{\text{env}}}, \end{aligned}$$

where

$$\tilde{\rho} = \sum_{\substack{E \in \text{sp } H_p \\ E' \in \text{sp } H_{\text{env}}}} P_{p,E} \rho_p P_{p,E} \otimes P_{\text{env},E'} \rho_\beta^{\otimes M} P_{\text{env},E'},$$

and $P_{p,E}, P_{\text{env},E'}$ are the spectral projections of H_p and H_{env} . Since $\rho_\beta^{\otimes M}$ commutes with H_{env} , this reduces to $\tilde{\rho} = \tilde{\rho}_p \otimes \rho_\beta^{\otimes M}$ with $\tilde{\rho}_p = \sum_{E \in \text{sp } H_p} P_{p,E} \rho_p P_{p,E}$. Hence, invoking (2.17), we obtain

$$\begin{aligned}
& g_n(\alpha_p, \alpha_{\text{env}}) \\
&= \log \text{Tr } U(n\tau, 0) (\tilde{\rho}_p e^{-\alpha_p \beta^* H_p} \otimes \rho_\beta^{\otimes M} e^{\alpha_{\text{env}} \beta H_{\text{env}}}) U(n\tau, 0)^* (e^{\alpha_p \beta^* H_p} \otimes e^{-\alpha_{\text{env}} \beta H_{\text{env}}}) \\
&= \log \text{Tr } (e^{\alpha_p \beta^* H_p} \otimes [\rho_\beta^{\alpha_{\text{env}}}]^{\otimes M}) U(n\tau, 0) (\tilde{\rho}_p e^{-\alpha_p \beta^* H_p} \otimes [\rho_\beta^{1-\alpha_{\text{env}}}]^{\otimes M}) U(n\tau, 0)^* \\
&= \log \text{Tr}_{\mathcal{H}_p} e^{\alpha_p \beta^* H_p} \mathcal{L}_{\alpha_{\text{env}}}^n (\tilde{\rho}_p e^{-\alpha_p \beta^* H_p}) \\
&= \log \text{Tr}_{\mathcal{H}_p} \mathcal{U}^{*n} (e^{\alpha_p \beta^* H_p}) \tilde{\mathcal{L}}_{\alpha_{\text{env}}}^n (\tilde{\rho}_p e^{-\alpha_p \beta^* H_p}) \\
&= \log \text{Tr}_{\mathcal{H}_p} e^{\alpha_p \beta^* H_p / 2} \tilde{\mathcal{L}}_{\alpha_{\text{env}}}^n (e^{-\alpha_p \beta^* H_p / 2} \tilde{\rho}_p e^{-\alpha_p \beta^* H_p / 2}) e^{\alpha_p \beta^* H_p / 2}.
\end{aligned}$$

Again, the number M of interactions is now irrelevant and we may consider arbitrary values of n . The commutation relations (2.2) imply $e^{\eta H_p} T e^{-\eta H_p} = e^{-\eta F} T$ so that, by (5.5),

$$\begin{aligned}
& e^{\alpha_p \beta^* H_p / 2} \tilde{\mathcal{L}}_{\alpha_{\text{env}}}^n (e^{-\alpha_p \beta^* H_p / 2} A e^{-\alpha_p \beta^* H_p / 2}) e^{\alpha_p \beta^* H_p / 2} \\
&= e^{(\alpha_p + \alpha_{\text{env}}) \beta E} p_- T^* A T + p_0 A + e^{-(\alpha_p + \alpha_{\text{env}}) \beta E} T A T^* \\
&= \tilde{\mathcal{L}}_{\alpha_p + \alpha_{\text{env}}}^n (A).
\end{aligned}$$

It follows that

$$g_n(\alpha_p, \alpha_{\text{env}}) = \log \text{Tr } I \tilde{\mathcal{L}}_{\alpha_p + \alpha_{\text{env}}}^n (\tilde{\rho}_p) = \log \text{Tr } \tilde{\rho}_p \tilde{\mathcal{L}}_{\alpha_p + \alpha_{\text{env}}}^{*n} (I) = n \log \theta(\alpha_p + \alpha_{\text{env}}),$$

from which we conclude that

$$\log \mathbb{E}^n [e^{\alpha(\Delta S_{p,n} - \Delta S_{\text{env},n})}] = n \log \theta(0) = 0,$$

which proves Part 1, from which Part 2,

$$g_n(\alpha) = \log \mathbb{E}^n [e^{\alpha \Delta S_{p,n}}] = \log \text{Tr } I \tilde{\mathcal{L}}_{\alpha}^n (\tilde{\rho}_p) = n \log \theta(\alpha), \quad (7.3)$$

immediately follows. Differentiation of the last identity at $\alpha = 0$ gives Part 3. Since $\theta(\alpha)$ is an entire function of α such that $\theta(0) = 1$, $\log \theta(\alpha)$ is an analytic function of α in a complex neighborhood of 0 and Part 4 follows from the Bryc theorem (see [Bry]). The Gärtner-Ellis theorem directly applies to give Part 5. Finally, the transient fluctuation theorem Part 6 is a direct consequence of the symmetry $\theta(1 - \alpha) = \theta(\alpha)$ (and hence of time reversal invariance) which implies

$$\begin{aligned}
\log \sum_s \mathbb{P}^n [\Delta S_{p,n} = -s] e^{-s} e^{\alpha s} &= \log \sum_s \mathbb{P}^n [\Delta S_{p,n} = s] e^{(1-\alpha)s} \\
&= g_n(1 - \alpha) = g_n(\alpha) = \log \sum_s \mathbb{P}^n [\Delta S_{p,n} = s] e^{\alpha s},
\end{aligned}$$

and hence $\mathbb{P}^n [\Delta S_{p,n} = -s] e^{-s} = \mathbb{P}^n [\Delta S_{p,n} = s]$.

7.2 Proof of Theorem 3.3

We now consider the case where $A = X$. The cumulant generating function of the increment ΔX_n is given by Equ. (7.2),

$$g_n(\eta) = \log \mathbb{Q}^n [e^{\eta \Delta X_n}] = \log \text{Tr} \mathcal{L}_0^n(\tilde{\rho}_p e^{-\eta X}) e^{\eta X}.$$

Using the factorization $\mathcal{L}_0^n = \mathcal{U}^n \circ \tilde{\mathcal{L}}_0^n$, we further get

$$\begin{aligned} g_n(\eta) &= \log \text{Tr} \tilde{\mathcal{L}}_0^n(e^{-\eta X/2} \tilde{\rho}_p e^{-\eta X/2}) \mathcal{U}^{*n}(e^{\eta X}) \\ &= \log \text{Tr}(e^{\eta X/2} \tilde{\mathcal{L}}_0^n(e^{-\eta X/2} \tilde{\rho}_p e^{-\eta X/2}) e^{\eta X/2})(e^{-\eta X/2} e^{in\tau H_p} e^{\eta X} e^{-in\tau H_p} e^{-\eta X/2}). \end{aligned}$$

We note that $e^{\eta X} T e^{-\eta X} = e^{\eta T}$ and $e^{\eta X} T^* e^{-\eta X} = e^{-\eta T^*}$ so that, by (5.5),

$$e^{\eta X/2} \tilde{\mathcal{L}}_0^n(e^{-\eta X/2} A e^{-\eta X/2}) e^{\eta X/2} = e^{-\eta} p_- T^* A T + p_0 A + e^{\eta} T A T^* = \tilde{\mathcal{L}}_{-\eta/\beta E}^n(A).$$

It follows that

$$g_n(\eta) = \log \text{Tr} \tilde{\mathcal{L}}_{-\eta/\beta E}^n(\tilde{\rho}_p)(e^{-\eta X/2} e^{in\tau H_p} e^{\eta X} e^{-in\tau H_p} e^{-\eta X/2}).$$

Part 2 of Lemma 6.1 yields the estimates

$$\log C_\eta^{-1} \text{Tr} \tilde{\mathcal{L}}_{-\eta/\beta E}^n(\tilde{\rho}_p) I \leq g_n(\eta) \leq \log C_\eta \text{Tr} \tilde{\mathcal{L}}_{-\eta/\beta E}^n(\tilde{\rho}_p) I,$$

and since $\text{Tr} \tilde{\mathcal{L}}_{-\eta/\beta E}^n(\tilde{\rho}_p) I = \text{Tr} \tilde{\rho}_p \tilde{\mathcal{L}}_{-\eta/\beta E}^{*n}(I) = \theta(-\eta/\beta E)^n$, we finally get

$$\frac{1}{n} g_n(\eta) = \log \theta \left(-\frac{\eta}{\beta E} \right) + O \left(\frac{1}{n} \right),$$

so that

$$g(\eta) = \lim_{n \rightarrow \infty} \frac{1}{n} g_n(\eta) = \log \theta \left(-\frac{\eta}{\beta E} \right),$$

which proves Part 1. Part 2 and Part 4 follow from the Gärtner-Ellis theorem while Part 3 follows from the Bryc theorem. Finally, the LDP implies

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\mathbb{Q}^n \left[\frac{\Delta X_n}{n} \in [-q - \delta, -q + \delta] \right]}{\mathbb{Q}^n \left[\frac{\Delta X_n}{n} \in [q - \delta, q + \delta] \right]} = - \inf_{|x-q| \leq \delta} I(-x) + \inf_{|x-q| \leq \delta} I(x)$$

and the fact that $I(-x) = I(x) + \beta E x$ leads to Part 5.

Acknowledgments This work was supported in part by the ANR contract HAMMARK (ANR-09-BLAN-0098-01) and by the Ministry of Higher Education and Research, Nord-Pas de Calais Regional Council and FEDER through the Contrat de Projets État Région (CPER) 2007-2013.

References

- [AM] Aschcroft, N.W. and Mermin, N.D.: *Solid State Physics*. Holt-Saunders International Editions (1976).
- [AP] Attal, S. and Pautrat, Y.: From repeated to continuous quantum interactions. *Ann. Henri Poincaré* **7**, 59–104 (2006).
- [ABGK] Avron, J.E., Bachmann, S., Graf, G.M. and Klich, I: Fredholm determinants and the statistics of charge transport. *Commun. Math. Phys.* **280**, 807–829 (2008).
- [B] Bellissard, J.: Coherent and dissipative transport in aperiodic solids. In *Dynamics of Dissipation*, P. Garbaczewski, R. Olkiewicz (Eds.). *Lecture Notes in Physics* **597**, 413–486 (2003).
- [BJM] Bruneau, L., Joye, A. and Merkli, M.: Asymptotics of repeated interaction quantum systems. *J. Funct. Anal.* **239**, 310–344 (2006).
- [BP] Bruneau, L. and Pillet, C.-A.: Thermal relaxation of a QED cavity. *J. Stat. Phys.* **134**, 1071–1095 (2009).
- [Bry] Bryc, W.: A remark on the connection between the large deviation principle and the central limit theorem. *Stat. Prob. Lett.* **18**, 253–256 (1993).
- [CDG] Cohen-Tannoudji, C., Dupont-Roc, J. and Grinberg, G.: *Atom–Photon Interactions*. Wiley, New York (1992).
- [CDM] Clark, J., De Roeck, W. and Maes, C.: Diffusive behavior from a quantum master equation. *arXiv:0812.2858* (2009).
- [dR] de Roeck, W.: Large deviation generating function for currents in the Pauli-Fierz model. *Rev. Math. Phys.* **21**, 549–585 (2009).
- [DF] De Roeck, W. and Fröhlich, J.: Diffusion of a massive quantum particle coupled to a quasi-free thermal medium. *arXiv:0906.5178* (2009).
- [DFP] De Roeck W., Fröhlich, J. and Pizzo, A.: Quantum Brownian motion in a simple model system. *Commun. Math. Phys.*, **293**, 361–398 (2010).
- [Du] Dutra, S.M.: *Cavity Quantum Electrodynamics*. Wiley, New York (2005).
- [DZ] Dembo, A. and Zeitouni, O.: *Large Deviations Techniques and Applications*. Springer-Verlag, New York (1998).

- [JPR] Jakšić, V., Pillet, C.-A. and Rey-Bellet, L.: Entropic fluctuations in statistical mechanics I. Classical dynamical systems. To appear in *Nonlinearity*, arXiv:1009.3248 (2010).
- [FZ] Fisher, P. A. and Zwerger, W.: Quantum Brownian motion in a periodic potential. *Phys. Rev. B* **32**, 6190–6206 (1985).
- [OLBC] Olver, W.J., Lozier, D.W., Boisvert, R.F. and Clark, C.W.: *NIST Handbook of Mathematical Functions*. Cambridge University Press (2010).
- [RM] Rondoni, R. and Mejía-Monasterio, C.: Fluctuations in nonequilibrium statistical mechanics: models, mathematical theory, physical mechanisms. *Nonlinearity* **20**, R1–R37 (2007).
- [Sch] Schrader, R.: Perron-Frobenius theory for positive maps on trace ideals. *Fields Inst. Commun.* **30** (2001).